arXiv:1610.05940v1 [cond-mat.quant-gas] 19 Oct 2016

Generalized Stefan-Boltzmann law

Gilles Montambaux
Laboratoire de Physique des Solides, CNRS, Université Paris-Sud, Université Paris-Saclay, 91405- Orsay
(Dated: October 20, 2016)

We reconsider the thermodynamic derivation by L. Boltzmann of the Stefan law and we generalize it for various different physical systems *whose chemical potential vanishes*. Being only based on classical arguments, therefore independent of the quantum statistics, this derivation applies as well to the saturated Bose gas in various geometries as to "compensated" Fermi gas near a neutrality point, such as a gas of Weyl Fermions. It unifies in the same framework the thermodynamics of many different bosonic or fermionic non-interacting gases which were until now described in completely different contexts.

Introduction — In 1879, J. Stefan found experimentally that the total power emitted by a black body scales as the fourth power of the temperature.[1] This law was derived theoretically by L. Boltzmann in 1884,[2] without any knowledge on the *spectral repartition* of the black body emission which was discovered later by M. Planck.[3] Nowadays, the now well-established expression of the Stefan-Boltzmann law with its correct prefactor[4]

$$u(T) = \frac{\pi^2 k_B^4}{15\hbar^3 c^3}\, T^4 \ , \qquad (1)$$

where $u = U/V$ is the energy per volume unit, is simply deduced in textbooks from the Planck law integrated over frequencies. It is however interesting to recall that the *total emission* can be derived from purely classical and thermodynamical grounds, as Boltzmann did. The main two ingredients for this derivation are the second law of thermodynamics and the result obtained by J.C. Maxwell[5] that the radiation exerts a pressure $p(T)$ proportional to the total energy density, $p(T) = \frac{1}{3}u(T)$. The 1/3 factor, obtained by Boltzmann, results from angular averaging in $3D$ and is indeed the inverse of the space dimensionality.

In this paper, we show how the Stefan-Boltzmann (SB) law can be generalized to any gas of free particles, and we stress the importance of a third ingredient for its derivation, a *zero chemical potential*, as it was implicit in the Boltzmann argument. In particular, we show how the simple relation between pressure and internal energy for a gas of massive particles $p(T) = \frac{2}{3}u(T)$ leads directly, *on purely classical grounds*, to the temperature dependence of the total energy of a $3D$ saturated Bose gas below the Bose-Einstein condensation temperature.

More generally, we consider free gases of particles with more general, including hybrid, dispersion relations, and show that the classical argument applies not only to the saturated Bose gas in various geometries (box or harmonic potentials), but also to gases of Fermions in various situations under the condition that the Fermi energy is temperature independent, which is the case for half-filled band systems. Such systems will be called "compensated Fermi gases". We regroup in the same framework the thermodynamic properties of many different bosonic or fermionic free gases, which until now were described in completely different contexts.

Stefan-Boltzmann law — We first briefly recall the derivation of this law as first theoretically derived by Boltzmann.[2] We start by considering the entropy $S(U, V)$. The variation of entropy for an infinitesimal transformation is

$$TdS = dU + pdV = \frac{\partial U}{\partial T}dT + \left(p + \frac{\partial U}{\partial V}\right) dV \ . \qquad (2)$$

Since the entropy is a total differential, the equality between cross derivatives leads to:

$$\frac{\partial p}{\partial T} = \frac{p+u}{T}. \qquad (3)$$

Inserting the Maxwell-Boltzmann relation $p(T) = u(T)/3$ in Eq.(3) gives immediately

$$\frac{1}{4}\frac{du}{u} = \frac{dT}{T} \ , \qquad (4)$$

leading to the Stefan law $u(T) = K_1 T^4$. Implicitly, in Eq.(2) the chemical potential is set to zero, which is essential for the demonstration here.

Of course the correct numerical factor can only be obtained from integration over frequencies of the spectral function $u(\nu, T)$ and requires the famous form of the Planck law discovered later in 1900.[3] However, the combination of universal constants could be obtained simply by dimensional arguments. The Planck constant was not introduced at that time, but the existence of a natural unit for action does not require quantization hypothesis. Indeed such natural unit was already introduced in 1899 by Planck[6](see supplementary material), so that Eq.(1) without the numerical factor can be obtained from purely classical arguments, h being a natural unit of action obtained experimentally.[6]

We now turn to the generalization of the SB law to a gas of massive particles with zero chemical potential.

Massive particles — Since the unique assumptions for deriving SB law are the thermodynamic principles and the relation between internal energy and pressure, it is

interesting to reconsider now the Boltzmann derivation in the case of massive particles. However an important remark is in order. Contrary to photons whose number is not fixed, the number of massive particles is in general fixed (in average) and controlled by a chemical potential μ. Eq.(2) should indeed be written as $TdS = dU + pdV - \mu dN$. Therefore, the use of Eq.(2) implies that the chemical potential is zero, which is actually the case for a gas of photons, leading to the SB law. If one now considers a gas of massive particles, we know from Maxwell-Boltzmann kinetic theory of gases that p and u are related by $p = \frac{2}{3}u$. The Boltzmann derivation still holds *under the condition that* $\mu = 0$. In this case, from (3) one gets, instead of (4):

$$\frac{2}{5}\frac{du}{u} = \frac{dT}{T} \tag{5}$$

so that $u(T) = K_2 T^{5/2}$. A dimensional argument which necessarily involves an action, the Boltzmann constant and a mass (instead of a velocity as before) leads to

$$u(T) \propto \frac{m^{3/2}}{h^3}(k_B T)^{5/2} \tag{6}$$

This behavior is actually well-known: this is the internal energy of the $3D$ saturated Bose gas or of a gas magnon excitations in a $3D$ ferromagnet. We stress that this expression does not require the knowledge of the thermal repartition of energy levels and no quantum mechanical input: the introduction of h as a unit of action in the prefactor does not imply any quantization. It is therefore quite interesting that the *thermodynamic of the saturated gas can be described in the framework of classical physics, with the unique hypothesis that* $\mu = 0$.

More generally, the Maxwell-Boltzmann kinetic theory of gases can be used for any dispersion relation between energy and momentum of the form $\epsilon \propto p^\beta$ in dimension d, and leads to the general relation $u(T) = \kappa p(T)$ with $\kappa = d/\beta$ ($\beta = 1$ for photons and $\beta = 2$ for massive particles). The thermodynamics of such gases is described below. .

General spectrum — Consider now any gas of particles at zero chemical potential whose relation between pressure and energy is given a linear relation $u(T) = \kappa p(T)$. Then Eqs.(4,5) have to be replaced by

$$\frac{1}{\kappa+1}\frac{du}{u} = \frac{dT}{T} \ . \tag{7}$$

Integration of this relation together with dimensionality arguments leads to the connection between energy-pressure relation and the generalized Stefan-Boltzmann law:

$$u(T) = \kappa p(T) \implies u(T) \propto \frac{(mc^2)^{d-\kappa}}{(hc)^d}(k_B T)^{\kappa+1} \tag{8}$$

which generalizes Eqs.(1,6).

We propose here a quicker, simpler and more general derivation : instead of pressure, which may not be well defined in certain situations that we will consider below, we consider the grand potential Ω, and suppose a relation of the form $U = -\kappa\Omega$ which generalizes the relation $u = \kappa p$. Then the thermodynamic relation between the grand potential and the energy:

$$U = \left.\frac{\partial \beta\Omega}{\partial \beta}\right|_\alpha \tag{9}$$

where $\alpha = \beta\mu$, together with $U = -\kappa\Omega$, leads immediately to

$$\left.\frac{\partial \beta U}{\beta U}\right|_\alpha = -\kappa\frac{\partial\beta}{\beta} \ . \tag{10}$$

Integration of this relation, combined with dimensional arguments, gives

$$U(T) = F(\alpha)\left(\frac{L}{hc}\right)^d (mc^2)^{d-\kappa}(k_B T)^{\kappa+1} \ , \tag{11}$$

where $F(\alpha)$ is a multiplicative integration function. Here no assumption is made on the structure of $F(\alpha)$ (which is actually an integral of Bose or Fermi factors). The generalized Boltzmann law (8) follows immediately when the chemical potential vanishes. Finally since the number of particles $N(T)$ is given by $N(T) = -\beta \left.\frac{\partial\Omega}{\partial\alpha}\right|_\beta$, we find that $U(T) \propto N(T) k_B T$.

For more complex dispersion relations as hybrid dispersion relations discussed later in sections, the calculation of pressure from kinetic theory may be difficult to achieve because of complex angular averages. The pressure may even not be properly defined. Then we may obtain the coefficient κ and deduce the SB law directly from the energy dependence of the density of states $\rho(\epsilon)$, when it is a power law. Here we rather consider the *integrated* density of states (also called counting function) denoted as $N_<(\epsilon)$. We start with an integrated density of states of the form $N_<(\epsilon) = B\epsilon^\kappa$, where B has necessarily the dimensions

$$B \propto L^d \frac{(mc^2)^{d-\kappa}}{(hc)^d} \ . \tag{12}$$

When $\alpha = 0$, the total energy $U(T)$ has the form

$$U(T) = \int \epsilon\rho(\epsilon) f(\beta\epsilon) d\epsilon \tag{13}$$

where no assumption is needed on the form on the function f except that it is a function of the single combination $\beta\epsilon$ (which is Wien's displacement law for light). This immediately leads to a power law $T^{\kappa+1}$ for the total energy and to the relation $U(T) = -\kappa\Omega(T)$ between

energy and grand potential. Therefore, we have the following correspondance

$$\rho(\epsilon) \propto \epsilon^{\kappa-1} \iff U = -\kappa\Omega \iff U \propto T^{\kappa+1} \ . \quad (14)$$

This is the main result of this paper. It may appear quite natural since the grand potential is actually a double integral of the density of states. It has however a profound origin which is the Wien's displacement law which is valid for any gas of *zero chemical potential*. Since it is obtained from purely classical arguments, no assumption is made on the nature of the particles. The argument applies as well as for Bosons or Fermions under the condition that $\mu = 0$. We now review many different physical systems whose thermodynamics is unified in this unique framework.

In the following, we introduce the thermal de Broglie wavelength $\lambda_{dB} = h/(2\pi m k_B T)^{1/2}$ and another characteristic wavelength for a massless spectrum $\lambda_W = hc/k_B T$, that we may call "Wien wavelength" since it separates classical and quantum behaviors. Then, the total number of particles in the saturated gas and the total energy can be rewritten as

$$N(T) \propto \frac{V}{\lambda_{dB}^{2(d-\kappa)} \lambda_W^{2\kappa-d}} \quad , \quad U(T) \propto N(T) k_B T \quad (15)$$

which is another form of the generalized SB law (8).

Hybrid spectrum — Up to now, we have only considered situations of either massless or massive particles, so that obviously a mass and a velocity cannot enter both in (8,15). One has respectively either $\kappa = d$ or $\kappa = d/2$, so that

$$N(T) = \frac{V}{\lambda_W^d} \qquad \text{or} \qquad N(T) = \frac{V}{\lambda_{dB}^d} \ . \quad (16)$$

Interesting is the more general situation of an *hybrid* spectrum, that we write in the short dimensionless notation

$$\epsilon \sim \sum_j A_j n_j^{\beta_j} \quad (17)$$

with $\beta_j = 2$ along d_m massive directions and $\beta_j = 1$ along d_c massless directions. A_j have the dimensions of an energy and n_j are positive integer numbers. The integrated density of states is still a power $N_<(\epsilon) = B\epsilon^\kappa$ with

$$\kappa = \sum_j \frac{1}{\beta_j} = d_c + \frac{d_m}{2} \quad (18)$$

and B is given by Eq.(12). Therefore the number of particles has the form

$$N(T) \propto \frac{V}{\lambda_{dB}^{d_m} \lambda_W^{d_c}} \quad (19)$$

and the expressions for energy and pressure result immediately.

Until now, we have only considered a free gas in a box of volume V. Particularly important in the context of Bose-Einstein condensation (BEC), is the case of a Bose gas trapped in an harmonic potential. To avoid any detailed calculations, it is useful to remark that the density of states of massive particles in an harmonic trap is the same as the one of free massless particles. This is of course reminiscent of the wave-particle duality that the radiation spectrum can be described either as an ensemble of resonators or as a gas of massless particles. The substitution to be done is

$$\frac{\pi c}{L} \to \omega \quad \text{or} \quad \frac{L}{\lambda_W} \to \frac{k_B T}{2\hbar\omega} \quad (20)$$

and we have $\kappa = d$. Therefore the integrated density of states has the form, for an isotropic harmonic well

$$N_<(\epsilon) \propto \left(\frac{\epsilon}{\hbar\omega}\right)^d \ . \quad (21)$$

Again $\hbar$ has been introduced from homogeneity, but the scaling dependence of the density of states does not need quantization. The above argument leads to

$$N(T) \propto \left(\frac{k_B T}{\hbar\omega}\right)^d \quad , \quad U(T) \propto N(T) k_B T \quad (22)$$

Notice that in this case Maxwell-Boltzmann argument cannot be used directly since volume and pressure are obviously not defined. However, Ref.11 defines an "harmonic volume" $\mathcal{V} = 1/\omega^3$, consistent with the above substitution (20), and the conjugated "harmonic pressure" $\mathcal{P} = -\partial\Omega/\partial\mathcal{V}$.[11]

For completion, we consider the hybrid situation with a box potential (or size L) profile in d_m directions and an harmonic profile in d_h direction. The substitution (20) in Eq.(19) gives

$$N(T) = \left(\frac{L}{\lambda_{dB}}\right)^{d_m} \left(\frac{k_B T}{\hbar\omega}\right)^{d_h} \ . \quad (23)$$

Compensated Fermi gas — Another situation where the Boltzmann derivation can be applied is the one of a Fermi gas, under the condition that $\mu = 0$. This peculiar situation occurs when the Fermi level is pinned at a neutrality point with electron-hole symmetry. This is for example the case for Dirac massless fermions, extensively studied to describe electronic states of graphene at half-filling, where the Fermi energy lies in the middle of the energy spectrum and the chemical potential *is independent of temperature*. We denote this system as a "compensated Fermi gas". The $T = 0K$ filled lower band plays the role of the condensate, and the finite T excited

electrons play the role of the saturated gas. This is however not a saturated gas in the sense that their number is proportional to the total number of Fermions (and not independent). As an example, the above arguments can be applied to the now recently extensively studied Weyl fermions in $3D$ semi-metals.[13]

Consider first the case of $2D$ Dirac fermions like in graphene, for which $d = 2$, $\beta = 1$, so that $\kappa = 2$ (the density of states is linear). The general relation (8) gives immediately

$$u(T) \propto \frac{(k_B T)^3}{(hc)^2} \tag{24}$$

leading to the well-known specific heat $\propto T^2$ of graphene for the half-filled band.[12] Similarly, in bilayer graphene, the spectrum is quadratic so that $\beta = 2$ and $\kappa = 1$, and from (8) we have

$$u(T) \propto \frac{m}{h^2}(k_B T)^2 \ . \tag{25}$$

Consider more generally Weyl Fermions in d dimensions. The energy spectrum being linear, one has $\beta = 1$, and $\kappa = d$, so that

$$u(T) \propto \frac{(k_B T)^{d+1}}{(hc)^d} \tag{26}$$

The thermodynamics of Weyl Fermions is therefore the same as the one of the black body.

Hybrid Fermions — It has been shown that under appropriate lattice distortion, the two Dirac points characteristic of the graphene electronic spectrum can merge into a single point, where the spectrum stays linear in one direction but becomes quadratic along the other direction.[14, 15] This hybrid point has been called a "semi-Dirac" point.[16] This is an interesting example of an hybrid spectrum characterized by both a mass and a velocity

$$\epsilon = \pm\sqrt{\left(\frac{p_x^2}{2m}\right)^2 + c^2 p_y^2} \tag{27}$$

whose thermodynamic properties have been studied.[14, 15] This is special case of a general dispersion relation of the form

$$\epsilon \sim \pm\sqrt{\sum_j A_j n_j^{2\beta_j}} \tag{28}$$

which leads to the same energy dependence of the density of states as a spectrum of the form (17) (see SM), and the energy density if given by Eq.(8) where $\kappa = d_c + d_m/2 = 3/2$. The thermodynamic properties of a semi-Dirac Fermi spectrum are therefore the same as the saturated Bose gas in a mixed boxed-harmonic potential. More generally a gas of Weyl fermions close to a merging point, the spectrum can transform from a full linear dispersion relation to a linear-linear-massive, or a linear-massive-massive, to a full massive spectrum. The corresponding thermodynamics is summarized in Table 1.

Conclusion — We have generalized the Boltzmann derivation of the Stefan-Boltzmann law to ideal gases of any kind of particles, under the condition that the chemical potential stays zero. From the Maxwell-Boltzmann relation between energy and pressure, one simply recovers the thermodynamic properties of a saturated gas of massive particle below the Bose-Einstein condensation temperature. Then we generalize the derivation to the case of hybrid dispersion relations, massive in some directions, massless in other directions. This situation is achieved in Bose gases confined in hybrid boxed-harmonic potentials. We have also shown that fermionic gases with particle-hole symmmetry behave indeed as saturated gases with fixed chemical potential. This remark leads to draw similarities between quite different problems like a semi-Dirac Fermi gas and a 2D Bose gas in a mixed boxed-harmonic potential. The recently studied physics of Weyl points in 3D is even richer since the dispersion relation can be linear in two directions and massive in a third direction, a situation thermodynamically similar to a 3D Bose gas confined in a harmonic trap in two directions and a boxed potential in the third direction. The unifying picture is summarized in Table 1.

The author acknowledges useful comments form J. Dalibard and J.-N. Fuchs.

.

	$\kappa = 1$	$\kappa = 3/2$	$\kappa = 2$	$\kappa = 5/2$	$\kappa = 3$
d=2	*massive fermions* $N(T) \propto \frac{L}{\lambda_{dB}^2}$	*semi-Dirac fermions* $N(T) \propto \frac{S}{\lambda_{dB}\lambda_W}$	black body *massless fermions* $N(T) \propto \frac{S}{\lambda_W^2}$		
d=2		BEC + $\sqcup$ + $\smile$ $N(T) \propto \frac{L}{\lambda_{dB}} \frac{k_B T}{\hbar\omega}$	BEC + $\smile$ + $\smile$ $N(T) \propto \left(\frac{k_B T}{\hbar\omega}\right)^2$		
d=3		F-magnons *massive fermions* $N(T) \propto \frac{V}{\lambda_{dB}^3}$	*hybrid MML fermions* $N(T) \propto \frac{V}{\lambda_{dB}^2 \lambda_W}$	*hybrid MLL fermions* $N(T) \propto \frac{V}{\lambda_{dB}\lambda_W^2}$	black body phonons, AF-magnons *massless fermions* $N(T) \propto \frac{V}{\lambda_W^3}$
d=3		BEC + $\sqcup$ + $\sqcup$ + $\sqcup$ $N(T) \propto \frac{V}{\lambda_{dB}^3}$	BEC + $\sqcup$ + $\sqcup$ + $\smile$ $N(T) \propto \frac{S}{\lambda_{dB}^2} \frac{k_B T}{\hbar\omega}$	BEC + $\sqcup$ + $\smile$ + $\smile$ $N(T) \propto \frac{L}{\lambda_{dB}} \left(\frac{k_B T}{\hbar\omega}\right)^2$	BEC + $\smile$ + $\smile$ + $\smile$ $N(T) \propto \left(\frac{k_B T}{\hbar\omega}\right)^3$
$u(T)$	T^2	$T^{5/2}$	T^3	$T^{7/2}$	T^4

Figure 1. This table summarizes the thermodynamic properties of various massive, massless or hybrid saturated ($\mu = 0$) gases. The properties depend both on the space dimensionality d and the proportionality coefficient κ between internal energy $u(T)$ and pressure $p(T)$. The symbols $\sqcup$ and $\smile$ denote respectively a box potential or a harmonic potential in a given direction. The examples of bosons are shown in normal characters, and the example of fermions are denoted in italics. MML denotes a massive-massive-massless spectrum, MLL a massive-linear-linear spectrum. One has $N(T) \propto T^\kappa$ and $U(T) \propto T^{\kappa+1}$.

Supplemental Material

PLANCK'S NATURAL UNIT OF ACTION

In his 1899 paper,[6] more than one year before his discovery of the black body law with quantization of energy exchange, M. Planck found a natural unit of action from his analysis of the Stefan-Boltzmann law and Wien's law:

$$u(\nu,T)=\frac{8\pi b}{c^3}\nu^3 e^{-a\nu/T} \tag{29}$$

where a and b are parameters to be fixed from the experiment. Comparing with experimental estimate of the total energy

$$u(T)=\int_0^\infty u(\nu,T)d\nu=\frac{48\pi b}{a^4c^3}\,T^4 \tag{30}$$

and Wien's displacement law, Planck could extract the parameters b/a^4 and a, and obtain the values $b = 6.885\,10^{-34}$ J.s and $a = 0.4818\,10^{-10}$ K.s. This was the first estimate of a unit of action, which was not yet related to any idea of quantization, and which will be denoted later as h in the 1900 discovery of the black body law. Moreover, in our modern notations, $a = h/k_B$ so that the obtained value of a leads to an estimate of $k_B = 1.43\,10^{-23}$J/K (that Planck will introduce one year later in his seminal paper).

EXACT THERMODYNAMIC RESULTS

We give here for completion the exact expressions for the density of states and thermodynamic quantities.

Integrated density of states

Here we give complete expressions fot the integrated density of states for particles with various dispersion relations.

• Free massive particles, $\epsilon=\frac{p^2}{2m}$:

$$N_<(\epsilon)=C_d\left(\frac{L}{2\pi}\right)^d\left(\frac{2m\epsilon}{\hbar^2}\right)^{d/2} . \tag{31}$$

where $C_d = \pi^{d/2}/\Gamma(1 + d/2)$ is the unit volume in d dimensions.

• Free massless particles, $\epsilon = |p|c$:

$$N_<(\epsilon)=A_d\left(\frac{L}{2\pi}\right)^d\left(\frac{\epsilon}{\hbar c}\right)^d . \tag{32}$$

• Particles in a isotropic harmonic potential, $\epsilon = (n_x + n_y + n_z)\hbar\omega$:

$$N_<(\epsilon)=\frac{1}{d!}\left(\frac{\epsilon}{\hbar\omega}\right)^d . \tag{33}$$

• $2D$ semi-Dirac fermions $\epsilon=\sqrt{\left(\frac{p_x^2}{2m}\right)^2+p_yc^2}$:[15]

$$N_<(\epsilon)=\left(\frac{L}{\hbar}\right)^2 C\frac{\sqrt{m}}{c}\epsilon^{3/2} \tag{34}$$

with $C=\frac{\Gamma(1/4)^2}{6\pi^{5/2}}$.

• $2D$ particles in a boxed-harmonic potential, $\epsilon=\frac{p_x^2}{2m}+n_y\hbar\omega$:

$$N_<(\epsilon)=\frac{2L}{3\pi}\frac{\sqrt{m}}{\hbar^2\omega}\epsilon^{3/2} . \tag{35}$$

• Particles with dispersion relation $\epsilon=\sum_j A_j n_j^{\beta_j}$, $n_j > 0$:

$$N_<(\epsilon)=\frac{\epsilon^\kappa}{\Gamma(1+\kappa)}\prod_j\frac{\Gamma(1+1/\beta_j)}{A_j^{1/\beta_j}} \tag{36}$$

with $\kappa = d_c + d_m/2$. All previous cases may be obtained form this formula with

$\beta_j = 1$, $A_i = \hbar\pi c_i/L_i$ or $A_i = \hbar\omega_i$

$\beta_j = 2$, $A_i = (\hbar^2\pi^2/(2m_iL_i^2)$, and the integrated density of states may be summarized in the form:

$$N_<(\epsilon)=\frac{\epsilon^\kappa}{\Gamma(1+\kappa)}\prod_{j=1}^{d_c}\left(\frac{L_j}{\pi\hbar c_j}\right)\prod_{j=1}^{d_m}\left(\sqrt{\frac{2\pi m}{h^2}}L_j\right) \tag{37}$$

and for an harmonic potential in d_c directions:

$$N_<(\epsilon)=\frac{\epsilon^\kappa}{\Gamma(1+\kappa)}\prod_{j=1}^{d_c}\left(\frac{1}{\hbar\omega_j}\right)\prod_{j=1}^{d_m}\left(\sqrt{\frac{2\pi m}{h^2}}L_j\right) . \tag{38}$$

• Particles with dispersion relation $\epsilon=\sqrt{\sum_j(A_jn_j)^{2\beta_j}}$, $n_j > 0$:

$$N_<(\epsilon)=\frac{\epsilon^\kappa}{\Gamma(1+\kappa/2)}\prod_j\frac{\Gamma(1+1/(2\beta_j))}{A_j^{1/\beta_j}} \tag{39}$$

With the above expressions of A_j, the integrated density of states can be summarized as:

$$N_<(\epsilon)=\frac{\epsilon^\kappa}{\Gamma(1+\kappa/2)}\Gamma(3/2)^{d_c}\Gamma(5/4)^{d_m} \times\prod_{j=1}^{d_c}\left(\frac{L_j}{\pi\hbar c_j}\right)\prod_{j=1}^{d_m}\left(\sqrt{\frac{2\pi m}{h^2}}L_j\right) \tag{40}$$

which is the same as for the dispersion $\epsilon=\sum_j A_j n_j^{\beta_j}$, within different numerical factors.

Bosons and Fermions with zero μ

Here we collect the exact thermodynamic quantities for the saturated Bose gas and the compensated Fermi gas ($\mu = 0$). We start from a power law for the integrated density of states , $N_<(\epsilon) = B\epsilon^\kappa$. The total energy is

$$U(T) = B\kappa \int_0^\infty \frac{\epsilon^\kappa}{e^{\beta\epsilon-\alpha} \pm 1} d\epsilon \ . \tag{41}$$

For the saturated Bose gas, the number of particles and the total energy read:[9]

$$\begin{aligned} N(T) &= B\Gamma(1+\kappa)\zeta(\kappa)\, T^\kappa \\ U(T) &= \kappa \frac{\zeta(\kappa+1)}{\zeta(\kappa)} N(T) k_B T \ . \end{aligned} \tag{42}$$

For the compensated Fermi gas, we have

$$\begin{aligned} N(T) &= B\Gamma(1+\kappa)(1-2^{1-\kappa})\zeta(\kappa)\, T^\kappa \\ U(T) &= \kappa \frac{2^\kappa - 1}{2^\kappa - 2} \frac{\zeta(\kappa+1)}{\zeta(\kappa)} N(T) k_B T \ . \end{aligned} \tag{43}$$

[1] J. Stefan, *Über die beziehung zwischen der wärmestrahlung und der temperatur*, Wiener Ber. II, **79**, 391 (1879)

[2] L. Boltzmann, *Ableitung des Stefan'schen Gesetzes, betreffend die Abhängigkeit der Wärmestrahlung von der Temperatur aus der electromagnetischen Lichttheorie*, Ann. der Phys. und Chemie. **22**, 291 (1884)

[3] M. Planck, *Über das gesetz der energieverteilung im normalspectrum*, Ann. d. Phys. **4**, 553 (1901)

[4] Also known as the Stefan-Boltzmann law is the epxression for the emitted power per unit area $\mathcal{P}(T) = \frac{c}{4}u(T) \equiv \sigma T^4$ where σ is named the Stefan-Boltzmann constant.

[5] J. C. Maxwell, *A treatise on electricity and magnetism*, Oxford (1873)

[6] M. Planck, *Über irreversible strahlungsvorgänge*, Sitz. der Preußischen Akademie der Wissenschaften, Berlin, **5**, 440 (1899)

[7] W. Wien, *Eine neue Beziehung der Strahlung schwarzer Körper zum zweiten Hauptsatz der Wärme theorie,* Sitz. der Preußischen Akademie der Wissenschaften, **1**, 55 (1893)

[8] W. Wien, *On the division of energy in the emission spectrum of a black body*, Phil. Mag. Series 5, vol. 43, **262**, 214 (1897)

[9] C. Pethick and H. Smith, *Bose-Einstein condensation in dilute gases*, Cambridge University Press (2002)

[10] A. Einstein, *Quantentheorie des Einatomigen Idealen Gases II*, Sitz. der Preußischen Akademie der Wissenschaften, pp. 3-14 (Berlin 1925)

[11] V. Romero-Rochín, *Equation of state of an interacting Bose gas confined by a harmonic trap: the role of the "harmonic" pressure*, Phys. Rev. Lett. **94**, 130601 (2005)

[12] See for example Y. N. Srivastava, A. Widom and J. Swain , *Thermodynamic Equations of State for Dirac and Majorana Fermions*, arXiv:hep-ph/9709434

[13] S. Rao, *Weyl semi-metals: a short review*, arXiv:1603.02821

[14] B. Wunsch, F. Guinea and F. Sols, *Dirac-point engineering and topological phase transitions in honeycomb optical lattices*, New J. Phys. **10**, 103027 (2008)

[15] G. Montambaux, F. Piéchon, J.-N. Fuchs and M.O. Goerbig, *A universal Hamiltonian for the motion and the merging of Dirac cones in a two-dimensional crystal* , Eur. Phys. J. B **72**, 509 (2009)

[16] S. Banerjee, R.R. Singh, V. Pardo and W.E. Pickett, *Tight-Binding Modeling and Low-Energy Behavior of the Semi-Dirac Point*, Phys. Rev. Lett. **103**, 016402 (2009)